

**SYNTHESIS AND LUMINESCENT PROPERTIES OF BIMETALLIC
GOLD(I) AND SILVER(I) PYRAZOLATE COMPLEXES**

NURUL HUSNA BINTI SABRAN

UNIVERSITI TEKNOLOGI MALAYSIA

SYNTHESIS AND LUMINESCENT PROPERTIES OF BIMETALLIC
GOLD(I) AND SILVER(I) PYRAZOLATE COMPLEXES

NURUL HUSNA BINTI SABRAN

A thesis submitted in fulfillment of the
requirements for the award of the degree of
Master of Science (Chemistry)

Faculty of Science
Universiti Teknologi Malaysia

FEBRUARY 2015

This work is dedicated to my beloved parent, Sabran and Sa'adiyah, and my siblings, who are always been there for me, and have never doubted my dreams, and also to all my friends, who has shared to joyful tears and get through the hard time together.

ACKNOWLEDGEMENT

First of all, all praise to Allah S.W.T., the Almighty for blessing and giving me inspiration to embark this work and strength to complete my master research as well as this thesis. This thesis would not have been possible without the guidance and the help of several individuals directly or indirectly and extended their valuable assistance in the preparation and completion of this study. I would like to express my gratitude to all who have helped me in the research and writing of this thesis.

Especially I would like to wish my greatest appreciation towards my supervisor, Dr. Hendrik Oktendy Lintang and my co-supervisors, Dr Leny Yulianti and Prof. Dr. Salasiah Endud for providing me opportunity to do this project on “*Synthesis and Luminescent Properties of Trinuclear Bimetallic Gold(I) and Silver(I) Pyrazolate Complexes*”. Without their guidance and encouragement, all of this work could not have been accomplished. I have gained a lot of knowledge and great experience during the project which is useful for my future and carrier.

This project also will not be successful without other people help and concern. Here in, I sincerely thanks to all staffs and my colleagues at Ibnu Sina Institute for Fundamental Science Studies and Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia especially my lab mates, Nur Fatihah Ghazalli, Juan MatMin, Mohamad Azani Abdul Kadir Jalani, and Abdul Hamid Umar whom rendered their help during the period of my project.

Last but not least, I would like express a sense of gratitude to my friends and my beloved parent of their moral support, strength, help and everything. Thank you very much.

ABSTRACT

Luminescent d^{10} complexes of gold(I) and silver(I) have particularly received much attention due to their phosphorescent characteristics originating from metal–metal interactions and their applications in organic light–emitting diodes, sensors, catalysis, optics, and photonics. While bimetallic gold(I) and silver(I) complexes have been synthesized as clusters or mixed–compounds, luminescent studies of molecular self–assembly of bimetallic gold(I) and silver(I) complexes have not yet been reported. Therefore, this research aimed to study the phosphorescent properties of molecularly self–assembled trinuclear bimetallic gold(I) and silver(I) pyrazolate complexes ($4[Au_3Pz_3]@[Ag_3Pz_3]R$) using fluorescence spectroscopy. Both trinuclear gold(I) and silver(I) pyrazolate complexes, $2[Au_3Pz_3]R$ and $3[Ag_3Pz_3]R$ were successfully synthesized from pyrazole ligands having different alkyl chains ($1(PzH)R$; $R = H, (OCH_3)_2Bn, (OC_{10}TEG)_3Bn$) with chloro(dimethylsulfide) gold(I) ($[Au(SMe_2)]Cl$) and silver(I) hexafluorophosphate ($AgPF_6$). Bimetallic pyrazolate complexes $4[Au_3Pz_3]@[Ag_3Pz_3]R$ were synthesized by stirring a mixture of $2[Au_3Pz_3]R$ and $3[Ag_3Pz_3]R$ in dry dichloromethane for 1 hour with molar ratios of $2[Au_3Pz_3]R$ to $3[Ag_3Pz_3]R$ of 1:1, 1:2, 1:3, 1:5, 1:10, 2:1, 3:1, and 5:1, whereas molar ratios of 1:1, 1:2, and 2:1 were used for synthesis of $(OC_{10}TEG)_3Bn$. At molar ratio of 1:1, the fluorescence spectrum of the resulting complex exhibited only one emission peak centered at 633 nm compared to 691 nm for $2[Au_3Pz_3]H$ and 471 nm for $3[Ag_3Pz_3]H$ when excited at 280 nm. Based on the luminescent changes at molar ratio 1:1, it is proposed that the formed bimetallic complex might be the gold(I)–silver(I) cluster, $4[Au_3Pz_3]@[Ag_3Pz_3]H$. On the other hand, the bimetallic pyrazolate complex obtained at molar ratio 1:1 was $4[Au_3Pz_3]@[Ag_3Pz_3](OCH_3)_2Bn$ when the alkyl chain was changed by $(OCH_3)_2Bn$. Two emission peaks at 463 and 606 nm were shown in fluorescence spectra where the intensity of the peak at 463 nm assigned to gold(I)–silver(I) interactions is relatively much lower in comparison to the peak at 606 nm of gold(I)–gold(I) interaction. The result obviously suggests molecular structural changes which may be associated to increase rigidity of side chain of the bimetallic complexes. When the alkyl chain was changed by $(OC_{10}TEG)_3Bn$, the resulting bimetallic amphiphilic complex, $4[Au_3Pz_3]@[Ag_3Pz_3](OC_{10}TEG)_3Bn$ with molar ratio 1:1 exhibited two emission peaks at 491 and 710 nm with almost the same intensity upon excitation at 276 nm, while $2[Au_3Pz_3](OC_{10}TEG)_3Bn$ and $3[Ag_3Pz_3](OC_{10}TEG)_3Bn$ showed emission peaks at 699 nm and 537 nm, corresponding to gold(I) and silver(I), respectively. These findings suggested the formation of bimetallic amphiphilic complex *via* self–assembly of alternating gold(I) and silver(I) complexes due to the more flexible amphiphilic alkyl chains. Of significance, the characteristic luminescent properties of $4[Au_3Pz_3]@[Ag_3Pz_3]R$ with different types of pyrazole ligands and molar ratios could be ascribed to changes of the gold(I)–silver(I) coordination in the self–assembled structures.

ABSTRAK

Kompleks pendarcahaya d^{10} aurum(I) dan argentum(I) telah menjadi tumpuan utama kerana ciri pendarfosfornya yang berpunca daripada interaksi antara logam-logam dan penggunaannya dalam diod pemancar cahaya organik, sensor, pemangkinan, optik dan fotonik. Walaupun kompleks dwilogam aurum(I) dan argentum(I) telah disintesis sebagai gugusan atau sebatian campuran, kajian pendarcahaya bagi molekul swahimpun kompleks dwilogam aurum(I) dan argentum(I) masih belum dilaporkan. Oleh itu, penyelidikan ini bertujuan untuk mengkaji ciri pendarfosfor molekul swahimpun kompleks trinuklear dwilogam aurum(I) dan argentum(I) pirazolat ($4[Au_3Pz_3]@[Ag_3Pz_3]R$) menggunakan spektroskopi fotopendarcahaya. Kedua-dua kompleks trinuklear aurum(I) dan argentum(I) pirazolat ($2[Au_3Pz_3]R$ dan $3[Ag_3Pz_3]R$) telah berjaya disintesis daripada ligan pirazola yang mempunyai rantai alkil yang berbeza ($R = H, (OCH_3)_2Bn, (OC_{10}TEG)_3Bn$) dengan aurum(I) kloro(dimetilsulfida) ($[Au(SMe_2)]Cl$) dan argentum(I) heksafluorofosfat ($AgPF_6$). Kompleks dwilogam pirazolat $4[Au_3Pz_3]@[Ag_3Pz_3]R$ telah disintesis dengan mencampurkan $2[Au_3Pz_3]R$ dan $3[Ag_3Pz_3]R$ dalam diklorometana kering selama satu jam dengan nisbah molar 1:1, 1:2, 1:3, 1:5, 1:10, 2:1, 3:1, dan 5:1, manakala nisbah molar 1:1, 1:2, dan 2:1 telah digunakan untuk sintesis $(OC_{10}TEG)_3Bn$. Pada nisbah molar 1:1, spektrum pendarfluor kompleks dwilogam yang terhasil mempamerkan satu puncak pancaran yang berpusat pada sekitar 633 nm berbanding 691 nm bagi $2[Au_3Pz_3]H$ dan 471 nm bagi $3[Ag_3Pz_3]H$ apabila diuja pada 280 nm. Berdasarkan perubahan pendarfluor pada nisbah molar 1:1, adalah dicadangkan kompleks dwilogam yang terbentuk berkemungkinan adalah gugusan aurum(I)–argentum(I), $4[Au_3Pz_3]@[Ag_3Pz_3]H$. Sebaliknya, kompleks dwilogam pirazolat yang diperolehi pada nisbah molar 1:1 adalah $4[Au_3Pz_3]@[Ag_3Pz_3](OCH_3)_2Bn$ apabila rantai alkil digantikan dengan $(OCH_3)_2Bn$. Dua puncak pancaran pada 463 dan 606 nm kelihatan dalam spektrum pendarfluor dengan keamatan puncak pada 463 nm yang dipadankan kepada interaksi aurum(I)–argentum(I) secara relatif lebih rendah berbanding keamatan puncak pada 606 nm hasil interaksi aurum(I)–aurum(I). Keputusan ini jelas mencadangkan perubahan struktur molekul mungkin disebabkan oleh peningkatan ketegaran rantai sisi kompleks dwilogam tersebut. Apabila rantai alkil digantikan dengan $(OC_{10}TEG)_3Bn$, kompleks amfifilik yang terhasil, $4[Au_3Pz_3]@[Ag_3Pz_3](OC_{10}TEG)_3Bn$ pada nisbah molar 1:1 menunjukkan dua jalur pancaran pada 491 dan 710 nm dengan keamatan puncak yang hampir sama apabila diuja pada 276 nm, manakala $2[Au_3Pz_3](OC_{10}TEG)_3Bn$ dan $3[Ag_3Pz_3](OC_{10}TEG)_3Bn$ menunjukkan jalur pancaran pada 699 and 537 nm, masing-masing sepadan dengan aurum(I) dan argentum(I). Keputusan ini mencadangkan pembentukan kompleks dwilogam amfifilik melalui penswahimpunan kompleks aurum(I) dan argentum(I) secara berselang-seli disebabkan rantai alkil amfifilik yang lebih fleksibel. Yang lebih penting, sifat ciri pendarcahaya $4[Au_3Pz_3]@[Ag_3Pz_3]R$ dengan jenis ligan pirazola dan nisbah molar yang berbeza boleh dianggap berpunca daripada perubahan pengkoordinatan aurum(I)–argentum(I) dalam struktur swahimpun tersebut.

TABLE OF CONTENT

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xii
	LIST OF FIGURES	xiv
	LIST OF ABBREVIATIONS	xix
	LIST OF APPENDICES	xx
1	INTRODUCTION	1
	1.1 Background of the Study	1
	1.2 Statement Problem	4
	1.3 Objectives of the Study	5
	1.4 Scope of the Study	6
	1.5 Significance of the Study	6
2	LITERATURE REVIEW	8
	2.1 Luminescence	8
	2.2 Luminescent Materials	10
	2.3 Luminescent of Transition Metal Complexes	13
	2.3.1 Transition d^{10} Metal Complexes	16
	2.3.1.1 Copper (I) Complexes	16

2.3.1.2 Silver (I) Complexes	18
2.3.1.3 Gold (I) Complexes	19
2.3.2 Transition Bimetallic of d^{10} Metal Complexes	21
2.3.2.1 Transition Bimetallic of d^{10} Metal Complexes with Different Group	21
2.3.2.2 Transition Bimetallic of d^{10} Metal Complexes with Same Group	23
2.4 Metallophilic Interaction	27
2.5 Self-Assembly	31
3 RESEARCH METHODOLOGY	33
3.1 General Instruments	33
3.2 Chemicals and Materials	34
3.3 Synthesis of Pyrazole Ligands (1 (PzH)R)	34
3.3.1 Synthesis of 1 (PzH)(OCH ₃) ₂ Bn (1b)	35
3.3.2 Synthesis of 1 (PzH)(OC ₁₀ TEG) ₃ Bn (1c)	36
3.4 Synthesis of Trinuclear Gold(I) and Silver(I) Pyrazolate Complexes (2 [Au ₃ Pz ₃]R and 3 [Ag ₃ Pz ₃]R)	37
3.4.1 Synthesis of Trinuclear Gold(I) Pyrazolate Complexes (2 [Au ₃ Pz ₃]R)	39
3.4.2 Synthesis of Trinuclear Silver(I) Pyrazolate Complexes (3 [Au ₃ Pz ₃]R)	41
3.5 Synthesis of Bimetallic Trinuclear Gold(I) and Silver(I) Pyrazolate Complexes (4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]R)	43
3.5.1 Synthesis of Bimetallic Gold(I) and Silver(I) Pyrazolate Complexes (4a , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]H)	44
3.5.2 Synthesis of Bimetallic Gold(I) and Silver(I) Pyrazolate Complexes (4b , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OCH ₃) ₂ Bn)	45

3.5.3	Synthesis of Bimetallic Gold(I) and Silver(I) Pyrazolate Complexes (4c , $4[\text{Au}_3\text{Pz}_3]@[\text{Ag}_3\text{Pz}_3](\text{OC}_{10}\text{TEG})_3\text{Bn}$)	46
3.6	Luminescent Properties of Bimetallic Gold(I) and Silver(I) Pyrazolate Complexes ($4[\text{Au}_3\text{Pz}_3]@[\text{Ag}_3\text{Pz}_3]\text{R}$)	47
4	RESULTS AND DISCUSSIONS	49
4.1	General Outline of the Research	49
4.2	Synthesis and Characterization of Pyrazole Ligands (1(PzH)R)	50
4.2.1	Synthesis and Characterization of 1(PzH)(OCH₃)₂Bn (1b)	50
4.2.2	Synthesis and Characterization of 1(PzH)(OC₁₀TEG)₃Bn (1c)	60
4.3	Synthesis and Characterization Trinuclear Gold(I) and Silver(I) Pyrazolate Complexes (2 $[\text{Au}_3\text{Pz}_3]\text{R}$ and 3 $[\text{Ag}_3\text{Pz}_3]\text{R}$)	69
4.3.1	Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (2 $[\text{Au}_3\text{Pz}_3]\text{R}$)	69
4.3.1.1	Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (2a , 2 $[\text{Au}_3\text{Pz}_3]\text{H}$)	70
4.3.1.2	Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (2b , 2 $[\text{Au}_3\text{Pz}_3](\text{OCH}_3)_2\text{Bn}$)	74
4.3.1.3	Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (2c , 2 $[\text{Au}_3\text{Pz}_3](\text{OC}_{10}\text{TEG})_3\text{Bn}$)	78
4.3.2	Synthesis and Characterization of Trinuclear Silver(I) Pyrazolate Complexes (3 $[\text{Ag}_3\text{Pz}_3]\text{R}$)	83

4.3.2.1 Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (3a , 3 [Ag ₃ Pz ₃]H)	84
4.3.2.2 Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (3b , 3 [Ag ₃ Pz ₃](OCH ₃) ₂ Bn)	86
4.3.2.3 Synthesis and Characterization of Trinuclear Gold(I) Pyrazolate Complexes (3c , 3 [Ag ₃ Pz ₃](OC ₁₀ TEG) ₃ Bn)	88
4.4 Synthesis and Characterization of Bimetallic Trinuclear Gold(I) and Silver(I) Pyrazolate Complexes (4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]R)	90
4.4.1 Synthesis and Characterization of Bimetallic Pyrazolate Complexes (4a , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]H)	91
4.4.2 Synthesis and Characterization of Bimetallic Pyrazolate Complexes (4b , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OCH ₃) ₂ Bn)	93
4.4.3 Synthesis and Characterization of Bimetallic Pyrazolate Complexes (4c , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OC ₁₀ TEG) ₃ Bn)	96
4.5 Luminescent Properties of Bimetallic Trinuclear Gold(I) and Silver(I) Pyrazolate Complexes (4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]R)	98
4.5.1 Bimetallic Pyrazolate Complexes (4a , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]H)	98
4.5.2 Bimetallic Pyrazolate Complexes (4b , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OCH ₃) ₂ Bn)	101
4.5.3 Bimetallic Pyrazolate Complexes (4c , 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OC ₁₀ TEG) ₃ Bn)	104

5	CONCLUSIONS AND RECOMMENDATIONS	108
	5.1 Conclusions	108
	5.2 Recommendations	111
	REFERENCES	112
	Appendices 1 – 37	124 – 160

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Luminescence phenomena and the methods of excitation	8
3.1	The molar ratio of 2a to 3a	45
3.2	The molar ratio of 2b to 3b	46
3.3	The molar ratio of 2c to 3c	47
4.1	¹ H-NMR data of 1b* and 1b**	52
4.2	¹³ C-NMR data of 1b* and 1b**	53
4.3	¹ H-NMR data of 1b** and 1b	56
4.4	¹³ C-NMR data of 1b** and 1b	57
4.5	¹ H-NMR data of 1c* and 1c**	61
4.6	¹³ C-NMR data of 1c* and 1c**	62
4.7	¹ H-NMR data of 1c** and 1c	64
4.8	¹³ C-NMR data of 1c** and 1c	66
4.9	¹ H-NMR data of 1a and 2a	70
4.10	¹³ C-NMR data of 1a and 2a	71
4.11	¹ H-NMR data of 1b and 2b	75
4.12	¹³ C-NMR data of 1b and 2b	76
4.13	¹ H-NMR data of 1c and 2c	79
4.14	¹³ C-NMR data of 1c and 2c	80
4.15	Mass and physical appearance of bimetallic pyrazolate complexes 4a	91
4.16	Mass and physical appearance of bimetallic pyrazolate complexes 4b	94
4.17	Mass and physical appearance of bimetallic pyrazolate complexes 4c	97

4.18	Luminescence values of λ_{max} of excitation and emission band, stokes shifts, and image under UV lamp of bimetallic pyrazolate complexes 4a	100
4.19	Luminescence values of λ_{max} of excitation and emission band, stokes shifts, and image under UV lamp of bimetallic pyrazolate complexes 4b	103
4.20	Luminescence values of λ_{max} of excitation and emission band, stokes shifts, and image under UV lamp of bimetallic pyrazolate complexes 4c	106

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Partial energy level diagram of photoluminescence molecule for fluorescence and phosphorescence phenomena	10
2.2	Representative framework structures of zeolites using faujasite type with compensation of negative charge	11
2.3	Emission of mixed lanthanide MOF at different temperature (10 and 300 K) when excited at 355 nm	12
2.4	Schematic structure of (Pd–TFFP) and (Ru–phen)	13
2.5	Schematic structure of ruthenium(II) bipyridine complex	13
2.6	Schematic structure of copper(I) alkynyl complexes	17
2.7	Molecular structure of a) mononuclear and b) binuclear of copper(I) 3,5-bis(trifluoromethyl) pyrazolate complexes	17
2.8	Photoluminescence spectra of trinuclear copper(I) pyrazolate complex versus temperature	18
2.9	Schematic of representing gold(I) and copper(I) pyrazolate complexes with different dendritic generations	21
2.10	Representative structure of heterometallic Au–Tl complex	22
2.11	Schematic representation of Pt_2Ag_4 and Pt_2Ag_2 complexes with their luminescence in the solid state upon exposure to UV lamp at an ambient temperature	22

2.12	Polymeric structure of bimetallic Au–Ag pentachlorophenyl linked by Au–Au short interactions	23
2.13	Polymeric structure of bimetallic Au–Ag pentafluorophenyl linked by Au–Au short interactions and emission spectrum at $\lambda_{\text{ex}} = 468 \text{ nm}$	24
2.14	Image of gold(I)–silver(I) cluster under daylight and under UV lamp when grinding and adding solvent	25
2.15	a) Schematic structure and image of nanocomposite of gold(I) pyrazolate complex as silver ions sensor, b) emission spectra of nanocomposite of gold(I) pyrazolate complex before and after dipping in a THF solution of silver ion	25
2.16	Representative structure of mixed–metal of gold(I) carbeniates and silver(I) 3,5–diphenylpyrazolates, 2:1 and 1:2	26
2.17	Representative structure of $[\text{Au}_3(\text{CH}_3\text{N}=\text{COCH}_3)_3]$ and the metallophilic interactions	28
2.18	Frontier orbital scheme for $[\text{Au}(\text{PR}_3)_3]$	28
2.19	X–ray crystal structure of “chair like” coordination geometry of $\{[3,5-(\text{CF}_3)_2\text{Pz}]\text{M}\}_3$ (M₃). M = Cu, Ag, and Au from left to right, respectively	29
2.20	Polymeric structure of gold(I) and silver(I) mixed–metal trinuclear complexes	30
2.21	Schematic representing of self–assembled structure metal(I) pyrazolate complexes	31
3.1	Synthetic scheme of pyrazole ligands (1(PzH)R) with different alkyl side chains from benzyl bromide (1*BnBr)	35
3.2	Synthetic scheme of trinuclear gold(I) (2[Au₃Pz₃]R) and silver(I) pyrazolate complexes (3[Ag₃Pz₃]R) from pyrazole ligands (1(PzH)R)	38

3.3	Synthetic scheme of trinuclear gold(I) ($2[\text{Au}_3\text{Pz}_3]\text{R}$) pyrazolate complexes from pyrazole ligands ($1(\text{PzH})\text{R}$)	39
3.4	Synthetic scheme of trinuclear silver(I) pyrazolate complexes ($3[\text{Ag}_3\text{Pz}_3]\text{R}$) from pyrazole ligands ($1(\text{PzH})\text{R}$)	41
3.5	Synthetic scheme of bimetallic trinuclear gold(I) and silver(I) pyrazolate complexes ($4[\text{Au}_3\text{Pz}_3]@[\text{Ag}_3\text{Pz}_3]\text{R}$)	44
3.6	Experimental set up for capturing image of bimetallic $4[\text{Au}_3\text{Pz}_3]@[\text{Ag}_3\text{Pz}_3]\text{R}$ pyrazolate complexes under UV lamp; a) front view and b) 90° side view	48
4.1	Schematic scheme of general research outline	50
4.2	Mechanism for the synthesis of $1b^{**}$ from $1b^*$	51
4.3	<i>Keto-enol</i> tautomerization	51
4.4	^1H -NMR spectra of a) $1b^*$ and b) $1b^{**}$	52
4.5	^{13}C -NMR spectra of a) $1b^*$ and b) $1b^{**}$	54
4.6	Schematic route of reaction mechanism of $1b$ from $1b^{**}$	55
4.7	^1H -NMR spectra of a) $1b^{**}$ and b) $1b$	56
4.8	^{13}C -NMR spectra of a) $1b^{**}$ and b) $1b$	58
4.9	Mass spectra of a) calculated and b) observed of $1b$	59
4.10	FT-IR spectrum of $1b$	60
4.11	^1H -NMR spectra of a) $1c^*$ and b) $1c^{**}$	61
4.12	^{13}C -NMR spectra of a) $1c^*$ and b) $1c^{**}$	63
4.13	^1H -NMR spectra of a) $1c^{**}$ and b) $1c$	65
4.14	^{13}C -NMR spectra of a) $1c^{**}$ and b) $1c$	67
4.15	Mass spectra of a) calculated and b) observed $1c$	68
4.16	FT-IR spectrum of $1c$	68
4.17	Mechanism reaction equation for the synthesis of $2[\text{Au}_3\text{Pz}_3]\text{R}$ from $1(\text{PzH})\text{R}$	69
4.18	^1H -NMR spectra of a) $1a$ and b) $2a$	71
4.19	^{13}C -NMR spectra of a) $1a$ and b) $2a$	72

4.20	Mass spectra of a) calculated and b) observed 2a	73
4.21	FT–IR spectra of 1a and 2a	73
4.22	Emission spectrum of 2a at $\lambda_{\text{ex}} = 280$ nm with its picture under UV lamp at wavelength of 254 nm	74
4.23	^1H –NMR spectra of a) 1b and b) 2b	75
4.24	^{13}C –NMR spectra of a) 1b and b) 2b	76
4.25	Mass spectra of a) calculated and b) observed 2b	77
4.26	FT–IR spectra of 1b and 2b	77
4.27	Emission spectrum of 2b at $\lambda_{\text{ex}} = 280$ nm with its picture under UV lamp at wavelength of 254 nm	78
4.28	^1H –NMR spectra of a) 1c and b) 2c	79
4.29	^{13}C –NMR spectra of a) 1c and b) 2c	81
4.30	Mass spectra of a) calculated and b) observed 2c	82
4.31	FT–IR spectrum of 2c	82
4.32	Emission spectrum of 2c at $\lambda_{\text{ex}} = 276$ nm with its picture under UV lamp at wavelength of 254 nm	83
4.33	Mechanism for the synthesis of (3 [Ag ₃ Pz ₃]R) from (1 (PzH)R)	84
4.34	Mass spectra of a) calculated and b) observed 3a	85
4.35	FT–IR spectra of 1a and 3a	85
4.36	Emission spectrum of 3a at $\lambda_{\text{ex}} = 280$ nm with its picture under UV lamp at wavelength of 254 nm	86
4.37	Mass spectra of a) calculated and b) observed 3b	87
4.38	FT–IR spectra of 1b and 3b	87
4.39	Emission spectrum of 3b at $\lambda_{\text{ex}} = 280$ nm with its picture under UV lamp at wavelength of 254 nm	88
4.40	Mass spectra of a) calculated and b) observed 3c	89
4.41	FT–IR spectra of 1c and 3c	89
4.42	Emission spectrum of 3c at $\lambda_{\text{ex}} = 276$ nm with its picture under UV lamp at wavelength of 254 nm	90
4.43	FT–IR spectra of bimetallic pyrazolate complexes 4a with different molar ratios of 2a to 3a	92
4.44	XRD diffractograms of bimetallic pyrazolate	93

	complexes 4a with different molar ratios of 2a to 3a	
4.45	FT–IR spectra of bimetallic pyrazolate complexes 4b with different molar ratios of 2b to 3b	95
4.46	XRD diffractograms of bimetallic pyrazolate complexes 4b with different molar ratios of 2b to 3b	96
4.47	XRD diffractograms of bimetallic pyrazolate complexes 4c with different molar ratios of 2c to 3c	97
4.48	Emission spectra of bimetallic pyrazolate complexes 4a with molar ratios 1:0, 0:1, 1:1, 1:2, 1:3, 1:5, 1:10, 2:1, 3:1, and 5:1 at 280 nm	99
4.49	Possible structure of bimetallic complex 4a ; 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃]H with molar ratio of 1:1 of 2a and 3a	101
4.50	Emission spectra of bimetallic pyrazolate complexes 4b with molar ratios 1:0, 0:1, 1:1, 1:2, 1:3, 1:5, 1:10, 2:1, 3:1, and 5:1 at 280 nm	102
4.51	Possible structure of bimetallic complex 4b ; 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OCH ₃) ₂ Bn with molar ratio of 1:1 of 2b and 3b	104
4.52	Emission spectra of bimetallic pyrazolate complexes 4c with molar ratios 1:0, 0:1, 1:1, 1:2, and 2:1 at 276 nm	105
4.53	Possible structure of bimetallic complex 4c ; 4 [Au ₃ Pz ₃]@[Ag ₃ Pz ₃](OC ₁₀ TEG) ₃ Bn with molar ratio of 1:1 of 2c and 3c	107

LIST OF ABBREVIATIONS

MOFs	- Metal Organic Frameworks
LF	- Ligand Field
MLCT	- Metal-to-Ligand Charge Transfer
IL	- Intraligand
MC	- Metal Centered
LMCT	- Ligand-to-Metal Charge Transfer
LMMCT	- Ligand-to-Metal-to-Metal Charge Transfer
VOCs	- Volatile Organic Compounds
NLO	- Non-linear Optic
OLEDs	- Organic Light-Emitting Devices
PL	- Photoluminescence
DNT	- 2,4-Dinitrotoluene
DMNB	- 2,3-Dimethyl-2,3-Dinitrobutane
UV	- Ultra Violet
DCM	- Dichloromethane
1*BnBr	- Benzyl Bromide
1**BnAcac	- Benzyl Acetyl Acetate
1(PzH)R	- Pyrazole Ligands
2[Au ₃ Pz ₃]R	- Trinuclear Gold(I) Pyrazolate Complexes
3[Ag ₃ Pz ₃]R	- Trinuclear Silver(I) Pyrazolate Complexes
4[Au ₃ Pz ₃]@[Ag ₃ Pz ₃]R	- Bimetallic Trinuclear Gold(I) and Silver(I) Pyrazolate Complexes
¹ H-NMR	- Proton Nuclear Magnetic Resonance
¹³ C-NMR	- Carbon Nuclear Magnetic Resonance
FT-IR	- Fourier Transform Infrared Spectroscopy
XRD	- X-ray diffractometer

LIST OF APPENDICES

APPENDIX NO.	TITLE	PAGE
1	Synthetic protocol for 1**BnAcac	124
2	¹ H-NMR spectrum of 1b** with chemical shift at -0.05–12.00 ppm	125
3	¹³ C-NMR spectrum of 1b** with chemical shift at 0–220 ppm	126
4	Mass spectrum of 1b**	127
5	Synthetic protocol for 1(PzH)R	128
6	¹ H-NMR spectrum of 1b with chemical shift at -0.05–12.00 ppm	129
7	¹³ C-NMR spectrum of 1b with chemical shift at 0–220 ppm	130
8	Mass spectrum of 1b	131
9	FT-IR spectra of 1b , 2b , and 3b	132
10	¹ H-NMR spectrum of 1c** with chemical shift at -0.05–12.00 ppm	133
11	¹³ C-NMR spectrum of 1c** with chemical shift at 0–220 ppm	134
12	Mass Spectrum of 1c**	135
13	¹ H-NMR spectrum of 1c with chemical shift at -0.05–12.00 ppm	136
14	¹³ C-NMR spectrum of 1c with chemical shift at 0–220 ppm	137
15	Mass spectrum of 1c	138
16	FT-IR spectra of 1c , 2c , and 3c	139

17	Synthetic protocol for 2a	140
18	Mass spectrum of 2a	141
19	FT-IR spectra of 2a and 3a	142
20	XRD diffractograms of 2a and 3a	143
21	Synthetic protocol for 2b	144
22	¹ H-NMR spectrum of 2b with chemical shift at -0.05–12.00 ppm	145
23	¹³ C-NMR spectrum of 2b with chemical shift at 0–220 ppm	146
24	Mass spectrum of 2b	147
25	XRD diffractograms of 2b and 3b	148
26	Synthetic protocol for 2c	149
27	¹ H-NMR spectrum of 2c with chemical shift at -0.05–12.00 ppm	150
28	¹³ C-NMR spectrum of 2c with chemical shift at 0–220 ppm	151
29	Mass spectrum of 2c	152
30	XRD diffractograms of 2c and 3c	153
31	Synthetic protocol for 3a	154
32	Mass spectrum of 3a	155
33	Synthetic protocol for 3b	156
34	Mass spectrum of 3b	157
35	Synthetic protocol for 3c	158
36	Mass spectrum of 3c	159
37	List of publications and conferences attended	160

CHAPTER 1

INTRODUCTION

1.1 Background of the Study

Luminescent materials are well-known as materials having phosphors or compounds that give emission properties when they absorb energy from light [1]. Since last few decades, many researchers have paid attention to develop the next generation of high performance luminescent materials in display, lighting, optical devices, sensing, and imaging [1–4]. However, some limitations on their physical properties such as quantum yield, spectral energy distribution, life time, and emission as well as their chemical stability and composition [1, 3] have been generally found as a parameter for reducing their performance.

Recently, researchers have been focused on development of new materials having high luminescent properties not only from organic or inorganic but also from both organic and inorganic phosphor compounds. Thereby, the development will potentially provide new functional luminescent materials such as metal organic phosphors [5], doped zeolites [6], metal organic frameworks (MOFs) [7], and composites [8]. For example, Lee *et al.* have reported that new blue emitting phosphor of $\text{Na}_x\text{Ca}_{1-x}\text{Al}_{2-x}\text{Si}_{2+x}\text{O}_8:\text{Eu}^{2+}$ (NCASO: Eu^{2+}) with excitation wavelength at wide spectral range from 250 to 420 nm [5] have been used as light emitting materials. Moreover, modification of Europium ion (Eu^{3+})–exchanged zeolite L with silylated β –diketone was reported to have a strong red emission due to an energy transfer of grafted molecules to the Eu^{3+} ions [6]. Recently, Cui *et al.* [7] have

reported that mix lanthanide ions with MOF showed two emission spectra at wide range temperature and were potentially used as a thermometer. In addition, composite luminescent materials containing of 5,10,15,20-tetrakis(2,3,4,5,6-pentafluorophenyl) porphyrin and tris(phenanthroline) with palladium(II) and ruthenium(I) as a metal ion (Pd-TFFP and Ru-phen) as reported previously in 2006 by Borisov and his co-workers can be used as dual oxygen and temperature sensor simultaneously [8].

Luminescent materials containing of metal organic complexes or known as organometallic have received great interest in recent years after the first report on photophysical and photochemical behaviors of ruthenium(II) bipyridine complexes by Adamson and Demas in 1971 [9]. Since this finding, numerous studies have been further developed and explored to significantly improve the performance. In 2001, Che and his co-workers have found that zinc(II) complex containing naphthyridyl ligand can give luminescent properties both in solution as an blue emission and in solid state as a white emission [10]. In some cases, transition metal complexes were getting more attention due to the capability to exhibit phosphorescent properties originating from triplet excited state of metal-to-ligand charge transfer (MLCT) [11–16]. For example, terpyridine ligand upon complexation with platinum ($[\text{Pt}(\text{terpy})\text{Cl}]^+$) was reported to form luminescence properties in both solid state and glass by Bailey and his group at 1995 [15]. However, the complexes were found to be non-emissive in the solution state due to a low energy of ligand field (LF) excited state from d–d transition.

Apart of monometallic complexes, some studies have highlighted that bimetallic or heterometallic complexes can be potentially used as optical devices [16], catalysts [17], and sensors [18]. Recently in 2013, cyclometalated d^8 platinum(II) complexes have been reported to self-assemble via both a weak non-covalent π – π and Pt–Pt interactions that are potentially used as organic light-emitting devices [19]. On following year, Li *et al.* was reported that bimetallic binuclear nickel and cobalt complexes of bis(benzotriazole iminophenolate) were highly reactive as a catalyst for copolymerization of cyclohexene oxide and carbon dioxide [17].

While most of studies were focused on d^6 and d^8 metal complexes, unique spectroscopy features of d^{10} metal complexes in their absorption and emission properties have been investigated based on a weak non-covalent bonding of metal-metal (metallophilic) interactions [20]. Recently, development on d^{10} metal complexes are not only from platinum group but also involving zinc(II), cadmium(II), gold(I), silver(I), and copper(I) complexes which are more considerable [21–24]. Peculiarly, gold(I), silver(I), and copper(I) complexes have received huge attention due to their characteristics of phosphorescent properties [25–32]. Gold(I), silver(I), and copper(I) pyrazolate complexes having different type of alkyl chains have been reported to self-assemble via weak metal-metal interactions [20] to form “chair like” coordination geometry of a cylindrical structure [29] and having high luminescent properties. In addition, Omary *et al.* have reported that dinuclear and mononuclear copper(I) and silver(I) complexes of 3,5-bis(trifluoromethyl)pyrazole ligand gave blue emission with short lifetimes [28]. Subsequently in 2005, this group has also reported to form supramolecular structure with luminescent changes upon complexation pyrazole ligands with gold(I), silver(I), and copper(I) metal ions [29]. Since these metal complexes have been reported to give high luminescent properties, it is interesting to investigate their potential applications such as vapochromic sensors where in 2011, trinuclear silver(I) pyrazolate complex was reported as a vapochromic selective sensor to benzene by Rawashdeh–Omary [32].

Instead of focusing on single metal complexes, some researchers also interested to study on the luminescence properties of bimetallic d^{10} complexes since these complexes showed great potential in various applications [33–35]. In 2004, heterometallic gold(I)–thallium(I) (Au–Tl) complex was reported as a vapochromic sensor due to its photophysical properties by Fernandez *et al.* [34]. On the other hands, Pina and his co-workers were reported that bimetallic gold–copper (Au–Cu) complex potentially can be used as a catalyst for oxidation of benzyl alcohol to benzaldehyde [35]. Nowadays, heterometallic gold–silver complexes have received much attention due to their photophysical and photochemical properties [36–43]. Pioneering of Omary and co-workers [36] and Burini and Fackler group [33, 37], the development of polymeric materials based on heterometallic gold–silver complexes

have been increased extensively [38] and showed potential applications in sensors [39], catalysts [40, 41], optics, and photonics [42]. For example, dimers of gold(I)–silver(I) mixed–metal complexes from gold(I) carbeniates, gold(I) benzylimidazoles, and silver(I) 3,5–diphenylpyrazoles were successfully synthesized by varying the molar ratios and found to be excellent candidate as a catalyst [40, 41].

Since investigation of bimetallic gold(I)–silver(I) complexes have been found to give excellent phosphorescent properties in the single crystals form [40], a lot of researches have been done to study their photochemical and photophysical properties. Recently in 2012, polynuclear organometallic of gold(I)–gold(I), gold(I)–silver(I), and gold(I)–copper(I) having bidentate ligands were reported to give high luminescent properties and can be used as active antimicrobial agents [44]. Although bimetallic gold(I)–silver(I) complexes have been synthesized in single crystals as a cluster or mix compounds, no example of luminescent studies for molecular assembly of bimetallic gold(I)–silver(I) complexes with various of alkyl side chains has yet been reported. Therefore, it would be a big challenge to study the phosphorescent properties of molecular assembled trinuclear bimetallic gold(I)–silver(I) pyrazolate complexes having different kinds of alkyl side chains and variation of the molar ratios by using photoluminescent spectroscopy. It is expected that the molar ratios of these bimetallic complexes will not only affect on the molecular structure of complexes but also possess unique phosphorescent properties.

1.2 Problem Statement

In 2003, Yang and Raptis have reported that trimeric gold(I) pyrazolate complex can self–assembly via gold–gold (aurophilic) interactions to show a red emission when excited at 230 nm [27]. On the other hands, the luminescent properties of trinuclear pyrazolate gold(I), silver(I), and copper(I) complexes have been further studied to investigate effect of the different metals on their supramolecular structure and as well as luminescent properties [29]. Since

developments on luminescent properties of single metal complexes have showed great potential in many applications, a few researches have also been done to study unique properties of bimetallic complexes.

Considering bimetallic gold–silver complexes have been found to give characteristic photophysical and photochemical properties for vapochemical sensors of volatile organic compounds (VOCs) [39], non–linear optic (NLO) materials [42], and organic light–emitting devices (OLEDs), it is an interesting challenge to synthesize molecular assembly of bimetallic gold(I)–silver(I) pyrazolate complexes having different kinds of alkyl side chains. However, no examples of self–assembled trinuclear bimetallic gold(I)–silver(I) pyrazolate complexes have been found to give phosphorescent properties with control of alkyl side chains. Thus, the goal of this research is to vary the molar ratios of mixed trinuclear gold(I) pyrazolate complex ($[\text{Au}_3\text{Pz}_3]$) and trinuclear silver(I) pyrazolate complex ($[\text{Ag}_3\text{Pz}_3]$) and then study the effect of the molar ratios on the luminescent properties. Finally, the resulting molecular assembled trinuclear bimetallic gold(I) and silver(I) pyrazolate ($[\text{Au}_3\text{Pz}_3]@[\text{Ag}_3\text{Pz}_3]$) complexes are expected to give unique phosphorescent properties.

1.3 Objectives of the Study

The objectives of this research can be separated as below:

- a) To synthesize pyrazole ligands with different alkyl side chains.
- b) To synthesize trinuclear gold(I) and silver(I)pyrazolate complexes having different alkyl side chains.
- c) To synthesize the trinuclear bimetallic gold(I) and silver(I) pyrazolate complexes by varying the molar ratios of gold(I)pyrazolate complex to silver(I)pyrazolate complex.
- d) To investigate the luminescent properties of the resulting trinuclear bimetallic gold(I) and silver(I)pyrazolate complexes.

1.4 Scope of the Study

The trinuclear gold(I) and silver(I) pyrazolate complexes were synthesized by using different pyrazole ligands attached with various alkyl chains; H, (OCH₃)₂Bn, and (OC₁₀TEG)₃Bn. The trinuclear bimetallic gold(I) and silver(I) pyrazolate complexes were prepared by mixing the gold(I) and silver(I) pyrazolate complexes with different molar ratios from 1:10, 1:5, 1:3, 1:2, 1:1, 1:2, 1:3, and 1:5, except for C₁₀TEG only using 1:2, 1:1, and 1:2.

The characterization of the resulting trinuclear gold(I), silver(I), and bimetallic gold(I) and silver(I) pyrazolate complexes was carried out by using Proton Nuclear Magnetic Resonance (¹H-NMR), Carbon Nuclear Magnetic Resonance (¹³C-NMR), Fourier Transform Infrared Spectroscopy (FT-IR), Ultraviolet Spectroscopy (UV), Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF-MS), and Liquid chromatography time-of-flight mass spectrometry (LC-TOF-MS). Structural analyses were confirmed by using X-Ray Diffraction (XRD), while the luminescent properties were studied by using spectrofluorophotometer. Photograph was taken by using digital camera with the macro zoom under UV lamp in the dark room.

1.5 Significance of the Study

The significance of this research is control the luminescent properties of bimetallic gold(I) and silver(I) pyrazolate complexes by synthesizing different alkyl side chains attached to pyrazole ligands. The resulting bimetallic complexes are expected to give unique luminescent properties by changing the molar ratios of gold(I) pyrazolate complex to silver(I) pyrazolate complex from 1:10, 1:5, 1:3, 1:2, 1:1, 1:2, 1:3, and 1:5, except for C₁₀TEG only using 1:2, 1:1, and 1:2. It is believed

that these bimetallic complexes with characteristics of luminescent properties can be used for potential applications in such as VOC sensors and NLO materials for imaging and optical data storage.

REFERENCE

1. Jüstel, T., Nikol, H., and Ronda, C. (1998). New Developments in the Field of Luminescent Materials for Lighting and Displays, *Angewandte Chemie International Edition*. **37**, 3084–3103.
2. Feldmann, C., Jüstel, T., Ronda, C. R., and Schmidt, P. J. (2003). Inorganic Luminescent Materials: 100 Years of Research and Application, *Advanced Functional Materials*. **13**, 511–516.
3. Blasse, G. (1989). New Luminescent Materials, *Chemistry of Materials*. **1**, 294–301.
4. Zhao, Q., Li, F., and Huang, C. (2010). Phosphorescent Chemosensors Based on Heavy–Metal Complexes, *Chemical Society Reviews*. **39**, 3007–3030.
5. Lee, G–Y, Han, J. Y, Im, W. B., Cheong, S. H., and Jeon, D. Y. (2012). Novel Blue–Emitting $\text{Na}_x\text{Ca}_{1-x}\text{Al}_{2-x}\text{Si}_{2+x}\text{O}_8:\text{Eu}^{2+}$ ($x = 0.34$) Phosphor with High Luminescent Efficiency for UV–Pumped Light–Emitting, *Inorganic Chemistry*. **51**, 10688–10694.
6. Li, H., Zhang, H., Wang, L., Mu, D., Qi, S., Hu, X., Zhang, L., and Yuan, J. (2012). Highly Luminescent Eu^{3+} –exchanged Zeolite L Crystals Resulting from Modification with Silylated β –diketone, *Journal of Materials Chemistry*. **22**, 9338–9342.
7. Cui, Y., Xu, H., Yue, Y., Guo, Z., Yu, J., Chen, Z., Gao, J., Yang, Y., Qian, G., and Chen, B. (2012). A Luminescent Mixed–Lanthanide Metal–Organic Framework Thermometer, *Journal of the American Chemical Society*. **134**, 3979–3982.

8. Borisov, S. M., Vasylevska, A. S., Krause, C., and Wolfbeis, O. S. (2006). Composite Luminescent Material for Dual Sensing of Oxygen and Temperature, *Advanced Functional Materials*. **16**, 1536–1542.
9. Adamson, A. W. and Demas, J. N. (1971). New Photosensitizer. Tris(2,2'-bipyridine)ruthenium(II) Chloride, *Journal of the American Chemical Society*. **93**, 1800–1801.
10. Che, C.-M., Wan, C.-W., Ho, K.-Y., and Zhou, Z.-Y. (2001). Strongly Luminescent Metal–Organic Compounds: Spectroscopic Properties and Crystal Structure of Substituted 1,8-naphthyridine and its Zinc(II) Complex, *New Journal of Chemistry*. **25**, 63–65.
11. Fernandez-M, V., Thorp-G, F. L., and Coogan, M. P. (2010). Application of d^6 Transition Metal Complexes in Fluorescence Cell Imaging, *Chemical Communications*. **46**, 186–202.
12. Kozhevnikov, V. N., Durrant, M. C., and Williams, J. A. G. (2011). Highly Luminescent Mixed-Metal Pt(II)/Ir(III) Complexes: Bis-cyclometallation of 4,6-Diphenylpyrimidine as a Versatile Route to Rigid Multimetallic Assemblies, *Inorganic Chemistry*. **50**, 6304–6313.
13. Chen, S.-C., Yu, R.-M., Zhao, Z.-G., Chen, S.-M., Zhang, Q.-S., Wu, X.-Y., Wang, F., and Lu, C.-Z. (2010). A Series of Polynuclear Complexes of d^{10} Metals with Interesting Luminescent Properties, *Journal of the American Chemical Society*. **10**, 1155–1160.
14. Yam, V. W.-W. and Lo, K. K.-W. (1999). *Multimetallic and Macromolecular Inorganic Photochemistry*. Marcel Dekker: New York. 31–112.
15. Bailey, J. M., Hill, M. G., Marsh, R. E., Miskowski, V. M., Schaefer, W. P., and Gray, H. B. (1994). Electronic Spectroscopy of Chloro (terpyridine) Platinum(II), *Inorganic Chemistry*. **34**, 4591–4599.

16. Chan, A. K.-W., Lam, E. S.-H., Tam, A. Y.-Y., Tsang, D. P.-K., Lam, W. H., Chan, M.-Y., Wong W.-T., and Yam, V. W.-W. (2013). Synthesis and Characterization of Luminescent Cyclometalated Platinum (II) Complexes of 1,3-Bis-Hetero-Azolybenzenes with Tunable Color for Applications in Organic Light-Emitting Devices through Extension of π Conjugation by Variation of the Heteroatom, *Chemistry-a European Journal*. **19**, 13910–13924.
17. Li, C.-H., Chuang, H.-J., Li, C.-Y., Ko, B.-T., and Lin, C.-H. (2014). Bimetallic Nickel and Cobalt Complexes as High Performance Catalysts for Copolymerization of Carbon Dioxide with Cyclohexene Oxide, *Polymer Chemistry*. **5**, 4875–4878.
18. Grate, J. W., Moore, L. K., Janzen, D. E., Veltkamp, D. J., Kaganove, S., Drew, S. M., and Mann, K. R. (2002). Steplike Response Behavior of a New Vapochromic Platinum Complex Observed with Simultaneous Acoustic Wave Sensor and Optical Reflectance Measurements, *Chemistry of Materials*. **14**, 1058–1066.
19. Diez, A., Fornies, J., Larraz, C., Lalinde, E., Lopez, J. A., Martin, A., Moreno, M. T., and Sicilia, V. (2010). Structural and Luminescence Studies on π - π and Pt-Pt Interactions in Mixed Chloro-Isocyanide Cyclometalated Platinum(II) Complexes, *Inorganic Chemistry*. **49**, 3239–3251.
20. Pyrkö, P. (1997). Strong Closed-Shell Interactions in Inorganic Chemistry, *Chemical Reviews*. **97**, 597–636.
21. Yam, V.W.W. and Lo, K.K.W. (1999). Luminescent Polynuclear d10 Metal Complexes, *Chemical Society Reviews*. **28**, 323–334.
22. Minghetti, G., Banditelli, G. and Bonati, F. (1979). Metal Derivatives of Azolate 3. The Pyrazolato Anion (and homologs) as a Mono- or Bidentate Ligand: Preparation and Reactivity of Tri-, Bi-, and Mononuclear Gold(I) Derivatives, *Inorganic Chemistry*. **18**, 658–663.

23. Ziolo, R. F., Lipton, S., and Dori, Z. J. (1970). The Photoluminescence of Phosphine Complexes of d^{10} Metals, *Journal of the American Chemical Society*. **1124**.
24. Kutal, C. (1990). Spectroscopic and Photochemical Properties of d^{10} Metal Complexes, *Coordination Chemistry Reviews*. **99**, 213–252.
25. Kim, S. J., Kang, S. H., Park K. M., Kim, H, Zin, W. C., Choi, M. G., and Kim, K. (1998). Trinuclear Gold(I) Pyrazolate Complexes Exhibiting Hexagonal Columnar Mesophases with Only Three Side Chains, *Chemistry of Materials*. **10**, 1889–1893.
26. Enomoto, M., Kishimura, A., and Aida, T. (2001). Coordination Metallacycles of an Achiral Dendron Self-Assemble via Metal–Metal Interaction to Form Luminescent Superhelical Fibers, *Journal of the American Chemical Society*. **123**, 5608–5609.
27. Yang, G and Raptis, R. G. (2003). Supramolecular Assembly of Trimeric Gold(I) Pyrazolate through Auophilic Attractions, *Inorganic Chemistry*. **42**, 261–263.
28. Omary, M. A., Rawashdeh-Omary, M. A., Diyabalanage, H. V. K., and Dias, H. V. R. (2003). Blue Phosphors of Dinuclear and Mononuclear Copper(I) and Silver(I) Complexes of 3,5-Bis(trifluoromethyl)pyrazolate and the Related Bis(pyrazolyl)borate, *Inorganic Chemistry*. **40**, 8612–8614.
29. Omary, M. A., Rawashdeh-Omary, M. A., Gonser, M. W. A., Elbjeirami, O., Grimes, T, and Cundari, T. R. (2005). Metal Effect on the Supramolecular Structure, Photophysics, and Acid–Base Character of Trinuclear Pyrazolate Coinage Metal Complexes, *Inorganic Chemistry*. **44**, 8200–8210.
30. Kishimura, A. (2005). Novel Luminescent Materials Based on the Self-Assembly via Metal–Metal Interactions among Group 11 Metal Ions, *Thesis Presented to the University of Tokyo*. 1–35.
31. Diaz, H. V. R. and Gamage, C. S. P. (2007). Arene-Sandwiched Silver(I) Pyrazolates, *Angewandte Chemie International Edition*. **46**, 2192–2194.

32. Rawashdeh-Omary, M. A., Rashdan, M. D., Dharanipathi, S., Elbjeirami, O., Ramesh, P., and Dias, H. V. R. (2011). On/off Luminescence Vapochromic Selective Sensing of Benzene and its Methylated Derivatives by a Trinuclear Silver(I) Pyrazolate Sensor, *Chemical Communications*. **47**, 1160–1162.
33. Burini, A., Bravi, R., Fackler, Jr J. P., Galassi, R., Grant, T. A., Omary, M. A., Pietroni, B. R., and Staples, R. J. (2000). Luminescent Chains Formed from Neutral, Triangular Gold Complexes Sandwiching Tl^{I} and Ag^{I} . Structures of $\{\text{Ag}([\text{Au}(\mu\text{-C}_2\text{N}_3\text{-bzim})]_3)_2\}\text{BF}_4\cdot\text{CH}_2\text{Cl}_2$, $\{\text{Tl}([\text{Au}(\mu\text{-C}_2\text{N}_3\text{-bzim})]_3)_2\}\text{PF}_6\cdot 0.5\text{THF}$ (bzim = 1-Benzylimidazolate), and $\{\text{Tl}([\text{Au}(\mu\text{-C}(\text{OEt})=\text{NC}_6\text{H}_4\text{CH}_3)]_3)_2\}\text{PF}_6\cdot\text{THF}$, with MAu_6 (M) Ag^+ , Tl^+) Cluster Cores, *Inorganic Chemistry*. **39**, 3158–3165.
34. Fernández, J.E., López-de-Luzuriaga, M.J., Monge, M., Olmos, E.M., Pérez, J., Laguna, A., Mohamed, A.A. and Fackler, Jr.P.J. (2004). $\{\text{Tl}[\text{Au}(\text{C}_6\text{Cl}_5)_2]\}_n$: A Vapochromic Complex, *Journal of the American Chemical Society*. **125**, 2022–2023.
35. Pina, C. D., Faletta, E., and Rossi, M. (2008). Highly Selective Oxidation of Benzyl Alcohol to Benzaldehyde Catalyzed by Bimetallic Gold–Copper Catalyst, *Journal of Catalysis*. **260**, 384–386.
36. Rawashdeh-Omary, M. A., Omary, M. A., and Fackler, Jr. J. P. (2002). Argento/Aurophilic Bonding in Organosulfur Complexes. The Molecular and Electronic Structures of the Heterobimetallic Complex $\text{AgAu}(\text{MTP})_2$, *Inorganica Chimica Acta*. **336**, 376–384.
37. Mohamed, M. A., Burini, A., and Fackler, Jr. J. P. (2005). Mixed–Metal Triangular Trinuclear Complexes: Dimers of Gold–Silver Mixed–Metal Complexes from Gold(I) Carbeniates and Silver(I) 3,5–Diphenylpyrazolates, *Journal of the American Chemical Society*. **127**, 5012–5013.
38. Laguna, A., Lasanta, T., Lopes-de-Luzuriaga, J. M., Monge, M., Naumov, P., and Olmos, M. E. (2010). Combining Aurophilic Interactions and Halogen

- Bonding to Control the Luminescence from Bimetallic Gold–Silver Clusters, *Journal of the American Chemical Society*. **132**, 456–457.
39. Fernández, E. J., Lopes–de–Luzuriaga, J. M., Monge, M., Olmos, M. E., Puelles, R. C., Laguna, A., Mohamed, A. A., and Fackler, Jr. J. P. (2008). Vapochromic Behavior of $\{\text{Ag}_2(\text{Et}_2\text{O})_2[\text{Au}(\text{C}_6\text{F}_5)_2]_2\}_n$ with Volatile Organic Compounds, *Inorganic Chemistry*. **47**, 8069–8076.
 40. Mohamed, A. A., Galassi, R., Papa, F., Burini, A., and Fackler, Jr. J. P. (2006). Gold(I) and Silver(I) Mixed-Metal Trinuclear Complexes: Dimeric Products from the Reaction of Gold(I) Carbeniates or Benzyylimidazoles with Silver(I) 3,5–Diphenylpyrazolate, *Inorganic Chemistry*. **45**, 7770–7776.
 41. Usón, R., Laguna, A., Laguna, M., and Manzano, B. R. (1984). Synthesis and Reactivity of Bimetallic Au–Ag Polyfluorophenyl Complexes; Crystal and Molecular Structures of $[\{\text{AuAg}(\text{C}_6\text{F}_5)_2(\text{SC}_4\text{H}_8)\}_n]$ and $[\{\text{AuAg}(\text{C}_6\text{F}_5)_2(\text{C}_6\text{H}_6)\}_n]$, *Journal Chemical Society, Dalton Transaction*. 285–292.
 42. Zhang, H., Zelmon, D. E., Deng, L., Liu, H.-K., and Teo, B. K. (2001). Optical Limiting Behavior of Nanosized Polycosahedral Gold–Silver Clusters Based on Third–Order Nonlinear Optical Effects, *Journal of the American Chemical Society*. **123**, 11300–11301.
 43. Usón, R., Laguna, A., Laguna, M., Manzano, B. R., and Tapia, A. (1985). Bimetallic Gold–Silver Pentachlorophenyl Complexes, *Inorganic Chimica Acta*. **101**, 151–153.
 44. Frik, M., Jimenez, J., Gracia, I., Falvello, L. R., Abi-Habib, S., Suriel, K., Muth, T. R., and Contel, M. (2012). Luminescent Di– and Polynuclear Organometallic Gold(I)–Metal (Au_2 , $\{\text{Au}_2\text{Ag}\}_n$ and $\{\text{Au}_2\text{Cu}\}_n$) Compounds Containing Bidentate Phosphanes as Active Antimicrobial Agents, *Chemistry–a European Journal*. **18**, 3659–3674.
 45. Valeur, B. (2001). *Molecular Fluorescence: Principles and Applications*. Wiley–VCH: Weinheim. 3–8.

46. Deshpande, S. S. (2001). Principles and Applications of Luminescence Spectroscopy, *Critical Reviews in Food Science and Nutrition*. **41**, 165.
47. Hashimoto, S. (2003). Zeolite Photochemistry: Impact of Zeolites on Photochemistry and Feedback from Photochemistry to Zeolite Science, *Journal of Photochemistry and Photobiology C: Photochemistry Review*. **4**, 19–49.
48. Valdes, M. G., Perez-Cordoves, A. I., and Diaz-Garcia, M. E. (2006). Zeolites and Zeolite-Based Materials in Analytical Chemistry, *Trend in Analytical Chemistry*. **25**, 24–30.
49. Kanan, S. M., Tripp, C. P., Austin, R. N., and Patterson, H. H. (2001). Photoluminescence and Raman Spectroscopy as Probes to Investigate Silver and Gold Dicyanide Clusters Doped in A-Zeolite and Their Photoassisted Degradation of Carbaryl, *Journal of Physical Chemistry*. **105**, 9441–9448.
50. Binnemans, K. (2009). Lanthanide-Based Luminescent Hybrid Materials, *Chemistry Review*. **109**, 4283–4374.
51. Jüstel, T., Wiechert, D. U., Lau, C., Sendor, D., and Kynast, U. (2001). Optically Functional Zeolites: Evaluation of UV and VUV Stimulated Photoluminescence Properties of Ce^{3+} - and Tb^{3+} -doped Zeolite X, *Advanced Functional Materials*. **11**, 105–110.
52. Allendorf, M. D., Bauer, C. A., Bhaktaa, R. K., and Houka, R. J. T. (2009). Luminescent Metal–Organic Frameworks, *Chemical Society Reviews*. **38**, 1330–1352.
53. Cui, Y., Yue, Y., Qian, G., and Chen, B. (2012). Luminescent Functional Metal–Organic Frameworks, *Chemical Reviews*. **112**, 1126–1162.
54. Kreno, L. E., Leong, K., Farha, O. K., Allendorf, M., Duyne, R. P. V., and Hupp, J. T. (2012). Metal–Organic Framework Materials as Chemical Sensors, *Chemical Reviews*. **112**, 1105–1125.

55. Chen, B., Wang, L., Zapata, F., Qian, G., Lobkovsky, E. B. (2008). A Luminescent Microporous Metal–Organic Framework for the Recognition and Sensing of Anions, *Journal of the American Chemical Society*. **130**, 6718–6719.
56. Lan, A., Li, K., Wu, H., Olson, D. H., Emge, T. J., Ki, W., Hong, M., and Li, J. (2009). A luminescent Microporous Metal–Organic Framework for the Fast and Reversible Detection of High Explosives, *Angewandte Chemie International Edition*. **48**, 2334–2338.
57. Crosby, G. A. (1975). Spectroscopic Investigations of Excited State of Transition–Metal Complexes, *Accounts of Chemical Research*. **8**, 231–238.
58. Kimachi, S., Ikeda, S., Miki, H., Azumi, T., and Crosby, G. A. (1994). Spectroscopic and Magnetic Studies of Complexes of d^{10} Closed Shell Ions, *Coordination Chemistry Reviews*. **132**, 43–50.
59. Crosby, G. A., Highland, R. G., and Truesdell K. A. (1985). Spectroscopic Properties of $(nd)^{10}$ Transition Metal Complexes, *Coordination Chemistry Reviews*. **64**, 41–52.
60. Liu, Q. D., Wang, R., and Wang S. (2004). Blue Phosphorescent Zn(II) and Orange Phosphorescent Pt(II) Complexes of 4,4′–diphenyl–6,6′–dimethyl–2,2′–bipyrimidine, *Journal Chemical Society, Dalton Transaction*. 2073–2079.
61. Nagle, K. J. (1988). $Tl_2Pt(CN)_4$: A Noncolumnar , Luminescent Form of $Pt(CN)_4^{2-}$ Containing Pt–Tl Bonds, *Journal of the American Chemical Society*. **110**, 319–321.
62. Passalacqua, R., Loiseau, F., Campagna, S., Fang, Y. Q., and Hanan G. S. (2003). In Search of Ruthenium(II) Complexes Based on Tridentate Polypyridine Ligands that Feature Long-lived Room-temperature Luminescence: The Multichromophore Approach, *Angewandte Chemie International Edition*. **115**, 1646–1649.
63. Lu, W., Mi, X. B., Chan, M. C. W., Hui, Z., Che, C. H., Zhu, N., and Lee, S. T. (2004). Light Emitting Tridentate Cyclometalated Platinum(II) Complexes

- Containing σ -Alkynyl Auxiliaries: Tuning of Photo- and Electrophosphorescence, *Journal of the American Chemical Society*. **126**, 4958–4971.
64. Wang, H. M. J., Chen, C. Y. L., and Lin, I. J. B. (1999). Synthesis, Structure, and Spectroscopic Properties Gold(I)–Carbene Complexes, *Organometallic*. **18**, 1216–1223.
65. Forward, J. M., Bohmann, D., Flacker, Jr. J. P., and Staples, R. J. (1995) Luminescence Studies of Gold(I) Thiolate Complexes, *Inorganic Chemistry*. **34**, 6330–6336.
66. Diez, J., Gamasa, M. P., Gimeno, J., Aguirre, A., and Garcia-Granda, S. (1991). $[\text{Cu}_3(\mu_3\text{-}\eta^1\text{-C}\equiv\text{CPh})_2(\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2)_3][\text{BF}_4]$, a Triangulo Copper(I) Complex with an Unprecedented Bicapping System of Two Asymmetric $\mu_3\text{-}\eta^1$ -Acetylide Ligands, *Organometallics*. **10**, 380–382
67. Yam, V. W. W., Lee, W. K., and Lai, T. F. (1993). Synthesis, Spectroscopy, and Electrochemistry of Trinuclear Copper(I) Acetylides. X-ray Crystal Structure of $[\text{Cu}_3(\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2)_3(\mu_3\text{-}\eta^1\text{-C}\equiv\text{C}^t\text{Bu})(\mu_3\text{-Cl})]\text{PF}_6$, *Organometallics*. **12**, 2383–2387.
68. Yam, V. W. W., Lee, W. K., Yeung, P. K. Y., and Philips, D. (1994). Detection of Transient $\text{Cu}^{\text{I}}\text{Cu}^{\text{I}}\text{Cu}^{\text{II}}$ Mixed-valence Acetylide Complexes from Nanosecond Transient Absorption Study, *Journal of Physical Chemistry*. **98**, 7545–7547.
69. Yam, V. W. W., Lee, W. K., Cheung, K. K., Crystall, B., and Philips, D. (1996). Synthesis, Structure, Photophysics, Time-resolved Emission Spectroscopy, and Electrochemistry of Luminescent Copper(I) Acetylide Complexes, *Journal Chemical Society, Dalton Transaction*. 3283–3287.
70. Lin, Y. Y., Lai, S. W., Che, C. M., Fu, W. F., Zhou, Z. Y., and Zhu, N. (2005). Structural Variations and Spectroscopic Properties of Luminescent Mono- and Multinuclear Silver(I) and Copper(I) Complexes Bearing Phosphine and Cyanide Ligands, *Inorganic Chemistry*. **44**, 1511–1524.

71. Yam, V. W. W., Fung, W. K. M., and Cheung, K. K. (1997). Synthesis, Photophysics, Crystal Structure of Hexanuclear Copper(I) and Silver(I) Acetylide Complexes, *Chemical Communications*. 963–964.
72. Lin, Y. Y., Lai, S. W., Che, C. M., Cheung, K. K., and Zhou, Z. Y. (2002). Luminescent Tetranuclear Silver(I) Arylacetylide Complexes Bearing Tricyclohexylphosphine Ligands: Synthesis, Molecular Structures, and Spectroscopic Comparison with Gold(I) and Copper(I) Arylacetalides, *Organometallics*. **21**, 2275–2282
73. Yam, V. W. W., Cheung, K. L., Yip, S. K., and Cheung, K. K. (2003). Synthesis, Characterization, Structure, and Luminescence Studies Mono-, Di-, and Trinuclear Gold Phosphine Alkynyl Complexes, *Journal of Organometallic Chemistry*. **681**, 196–209.
74. Zyl, W. E. V., Lopes-de-Luzuriaga, J. M., Mohamed, A. A., Staples, R. J., and Fackler, Jr. J. P. (2002). Dinuclear Gold(I) Dithiophosphonate Complexes: Synthesis, Luminescent Properties, and X-ray Crystal Structures of $[\text{AuS}_2\text{PR}(\text{OR}')]_2$ ($\text{R} = \text{Ph}$, $\text{R}' = \text{C}_5\text{H}_9$; $\text{R} = 4\text{-C}_6\text{H}_4\text{OMe}$, $\text{R}' = (1S,5R,2S)\text{-}(+)\text{-Menthyl}$; $\text{R} = \text{Fc}$, $\text{R}' = (\text{CH}_2)_2\text{O}(\text{CH}_2)_2\text{OMe}$), *Inorganic Chemistry*. **41**, 4579–4589.
75. Schmidbaur, H. (2000). The Auophilicity Phenomenon: A Decade of a Experimental Findings, Theoretical, Concepts, and Emerging Application, *Gold Bulletin*. **33**, 3–10.
76. Mansour, M. A., Connick, W. B., Lachicotte, R. J., Gysling, H. J., and Eisenberg, R. (1998). Linear Chain Au(I) Dimer Compounds as Environmental Sensors: A Luminescent Switch for the Detection of Volatile Organic Compounds, *Journal of the American Chemical Society*. **120**, 1329–1330.
77. Ito, H., Saito, T., Oshima, N., Kitamura, N., Ishizaka, S., Hinatsu, Y., Wakeshima, M., Kato, M., Tsuge, K., and Sawamura, M. (2008). Reversible Mechanochromic

- Luminescence of $[(C_6F_5Au)_2(\mu-1,4\text{-diisocyanobenzene})]$, *Journal of the American Chemical Society*. **130**, 10044–10045.
78. Lintang, H. O., Kinbara, K., Yamashita, T., and Aida, T. (2010). Self–Repair of a One–Dimensional Molecular Assembly in Mesoporous Silica by a Nanoscopic Template Effect, *Angewandte Chemie International Edition*. **49**, 4241–4245.
 79. Akatsu, S., Kanematsu, Y., Kurihara, T., Sueyoshi, S., Arikawa, Y., Onishi, M., Ishizaka, S., Kitamura, N., Nakao, Y., and Sakaki, S. (2012). Synthesis and Luminescent Properties of 3,5–Diphenylpyrazolato–Bridged Heteropolynuclear Platinum Complexes. The Influence of Chloride Ligands on the Emission Energy Revealed by the Systematic Replacement of Chloride Ligands by 3,5–Dimethylpyrazolate, *Inorganic Chemistry*. **51**, 7977–7992.
 80. Fernández, E. J., Laguna, A., Lopes–de–Luzuriaga, J. M., Monge, M., Montiel, M., Olmos M. E., and Rodríguez–Castillo, M. (2006). Photophysical and Theoretical Studies on Luminescent Tetranuclear Coinage Metal Building Blocks, *Organometallics*. **25**, 3639–3646.
 81. Ni, W.–X., Qiu, Y.–M., M. Li, Zheng, J., R. Sun W.–Y., Zhan, S.–Z., Ng, S. W., and Li, D. (2014). Metallophilicity–Driven Dynamic Aggregation of a Phosphorescent Gold(I) –Silver(I) Cluster Prepared by Solution–Based and Mechanochemical Approaches, *Journal of the American Chemical Society*. **136**, 9532–9535.
 82. Lintang, H. O., Kinbara, K., Yamashita, T., and Aida, T. (2012). Metal–Ion Permeation in Congested Nanochannels: The Exposure Effect of Ag^+ Ions on the Phosphorescence Properties of a Gold(I)–Pyrazolate Complex that is Confined in the Nanoscopic Channels of Mesoporous Silica, *Chemistry–an Asian Journal*. **7**, 2068–2072.
 83. Gade, L. H. (1997). Luminescent Gold(I) Compounds, *Angewandte Chemie International Edition*. **36**, 1171–1173.

84. Vickery, J. C., Olmstead, M. M., Fung, E. Y., and Balch, A. L. (1997). Solvent–Stimulated Luminescence from the Supramolecular Aggregation of a Trinuclear Gold(I) Complex that Displays Extensive Intermolecular Au–Au Interactions, *Angewandte Chemie International Edition*. **36**, 1179–1181.
85. Schmidbaur, H. (1995). High–Carat Gold Compounds, *Chemical Society Reviews*. 391–400.
86. Lasanta, T., Olmos, M. E., Laguna, A., Lopez, J. M., Naumov, P. (2011). Making the Golden Connection: Reversible Mechanochemical and Vapochemical Switching of Luminescence from Bimetallic Gold –Silver Cluster Associated through Aurophilic Interactions, *Journal of the American Chemical Society*. **133**, 16358–16361.
87. Kishimura, A., Yamashita, T., Yamaguchi, K., and Aida, T. (2005). Rewritable Phosphorescent Paper by the Control of Competing Kinetic and Thermodynamic Self–Assembling Events, *Nature Materials*. **4**, 546–549.